

Refractometry

Chapter 3 of Nesse

ERSC 2P22 – Brock University Greg Finn

Refractometry


Defined

- Requires having a set of immersion oils of _____ RI and comparing the mineral (unknown) to the oil

ERSC 2P22 – Brock University Greg Finn

Refraction of Light

- If $n_{oil} = n_{mineral}$, light passes through the oil-mineral boundary unrefracted
 - the grain does not stand out
- If $n_{oil} \neq n_{mineral}$, light passing through the oil-mineral boundary is refracted
 - the grain appears to stand out



ERSC 2P22 – Brock University Greg Finn

Relief

- Relief -

ERSC 2P22 – Brock University Greg Finn

Strong Relief

garnet

Mineral grain clearly stands out from the mounting material
 $RI_{\text{mineral}} - RI_{\text{mm}} > 0.12$ RI units

ERSC 2P22 – Brock University Greg Finn

Moderate Relief

apatite

$RI_{\text{mineral}} - RI_{\text{mm}}$ between 0.04 and 0.12 RI units

ERSC 2P22 – Brock University Greg Finn



Relief

- Minerals may exhibit:
 - High Relief
 - $RI_{\text{mineral}} > RI_{\text{oil}}$
 - eg. garnet, $RI = 1.76$
 - Low Relief
 - $RI_{\text{mineral}} < RI_{\text{oil}}$
 - eg. fluorite, $RI = 1.434$
- Useful to know whether the mineral RI is $>$ or $<$ oil - **BECKE LINE METHOD**

ERSC 2P22 - Brock University Greg Finn

Becke Line Method

- **Becke Line** -

ERSC 2P22 - Brock University Greg Finn

Becke Line Method

- Observing a Becke Line
 1. Use medium or high power
 2. Close down aperture diaphragm
 3. For high power flip in the auxillary condenser
- **Lower the stage, the Becke line will move into the material with the higher RI**

ERSC 2P22 - Brock University Greg Finn

Lens Effect

$n_{min} > n_{oil}$

Convergent Lens

Normal to Mineral Boundary

$n_{min} < n_{oil}$

Divergent Lens

The mineral concentrates the light into a convergent cone above the mineral. The light is refracted towards the normal on entering the mineral and away from the normal on exiting the mineral.

The mineral acts as a divergent lens and concentrates the light around the margins of the mineral. The light is refracted away from the normal on entering the mineral and towards the normal on exiting the mineral.

ERSC 2P22 - Brock University Greg Finn

Internal Reflection

$n_{min} > n_{oil}$

$n_{min} < n_{oil}$

Rays **1** and **4** are **refracted** into the mineral. Rays **2** and **3** are **internally reflected** because they strike the boundary at an angle greater than the critical angle. The Becke Line is formed by the concentration of light inside the mineral.

Rays **2** and **3** are **refracted** out of the mineral. Rays **1** and **4** are **internally reflected** because they strike the boundary at an angle greater than the critical angle. The Becke Line is formed by the concentration of light outside the mineral.

ERSC 2P22 - Brock University Greg Finn

Becke Line Movement

$n_{min} < n_{oil}$

The cone of light diverges above the mineral grain. Lowering the stage from F_1 to F_2 , the Becke line will appear to move out of the grain into the oil, i.e. into the material with the higher RI.

$n_{min} > n_{oil}$

The cone of light converges above the mineral grain. Lowering the stage from F_1 to F_2 , the Becke line will appear to move out of the oil into the grain, again into the material with the higher RI.

ERSC 2P22 - Brock University Greg Finn

Becke Line Movement

THE BECKE LINE WILL ALWAYS MOVE ONTO THE MATERIAL WITH THE HIGHER REFRACTIVE INDEX AS THE STAGE IS LOWERED

ERSC 2P22 - Brock University Greg Finn

RI in Thin Section

- Becke lines will form at the boundary between two materials
- Compare the unknown to a substance of known RI
 - Epoxy, $n = 1.54$
 - Quartz, $n = 1.544, n = 1.553$

ERSC 2P22 - Brock University Greg Finn
